# On the perimeter of a point set in the plane

Vasilis Capoyleas \*

János Pach †

## Abstract

Let  $\{c_0, \ldots, c_n\}, \{c'_0, \ldots, c'_n\}$  be two point sets in the plane satisfying  $|c_i - c_j| \le |c'_i - c'_j|$  for all i and j. By a theorem of Sudakov and Alexander, the perimeter of the convex hull of  $\{c_0, \ldots, c_n\}$  does not exceed the perimeter of the convex hull of  $\{c'_0, \ldots, c'_n\}$ . We give a simple proof of this result and establish a similar theorem in the case when the Euclidean distance is replaced by the maximum norm. We point out the close relationship between these questions and a longstanding open problem due to Thue Poulsen, Kneser and Hadwiger.

### 1 The union of balls.

More than 35 years ago E. Thue Poulsen [TP], M. Kneser [K] and H. Hadwiger [H] proposed the following conjecture which has attracted the interest of many geometers but is still open. Let  $\{C_0, \ldots, C_n\}, \{C'_0, \ldots, C'_n\}$  be two collections of disks of radius r in the plane. Let  $c_i$  and  $c'_i$  denote the center of  $C_i$  and  $C'_i$ , respectively, and assume that

$$|c_i - c_j| \le |c'_i - c'_j|$$
 for all  $i$  and  $j$ .

Then

$$\operatorname{Area}(\bigcup_{i=0}^{n} C_{i}) \leq \operatorname{Area}(\bigcup_{i=0}^{n} C'_{i}). \tag{1}$$

W. Habicht (see [K]) and B. Bollobás [Bo] settled the special case when the system  $\{C_0, \ldots, C_n\}$  can be continuously transformed into  $\{C'_0, \ldots, C'_n\}$  so that during the transformation the mutual distances between the centers do not decrease.

Assume now that (1) is true, and let the radii r of the disks tend to infinity. It is easy to see that

$$\operatorname{Area}(\bigcup_{i=0}^{n} C_{i}) = r^{2}\pi + r \operatorname{Per conv}\{c_{0}, \dots, c_{n}\}\$$

+Area conv
$$\{c_0,\ldots,c_n\}+O(\frac{1}{r}),$$

where Per and conv stand for the perimeter and the convex hull, respectively. Similarly,

Area
$$(\bigcup_{i=0}^{n} C'_i) = r^2 \pi + r \operatorname{Per conv}\{c'_0, \dots, c'_n\}$$

+Area conv
$$\{c'_0,\ldots,c'_n\}+O(\frac{1}{r}),$$

Now (1) immediately yields

$$\operatorname{Per conv}\{c_0,\ldots,c_n\} \leq \operatorname{Per conv}\{c'_0,\ldots,c'_n\}. \tag{2}$$

The aim of the present note is to give a simple elementary proof of this weaker assertion, which was first established by Sudakov ([S]) and rediscovered by Alexander ([A]). Our approach is similar to the one followed in Alexander's paper, but it avoids using Schläfli's formula. Both proofs are based on a simple property of simplices (Lemma 1 in the next section).

**Theorem 1** Let  $\{c_0, \ldots, c_n\}, \{c'_0, \ldots, c'_n\}$  be two point sets in the plane satisfying  $|c_i-c_j| \le |c'_i-c'_j|$  for all i and j. Then the perimeter of the convex hull of  $\{c_0, \ldots, c_n\}$  does not exceed the perimeter of the convex hull of  $\{c'_0, \ldots, c'_n\}$ .

Let us recall first some basic notions and results from the theory of convex bodies. Let  $B^n$  denote the n-dimensional unit ball, and let  $S^{n-1}$  be the boundary of  $B^n$ . For any convex set  $K \subseteq \mathbb{R}^n$ , let  $K + rB^n$  denote the parallel body of K with radius r, i.e., the set of all points of the space whose distance from at least one element of K is at most r. Let  $\operatorname{Vol}_n$  stand for the n-dimensional volume. It is well-known (see e.g. Bonnesen-Fenchel [BF], Busemann [Bu], Leichtweiss [L]) that  $\operatorname{Vol}_n(K + rB^n)$  can be expressed as a polynomial of degree n in r,

$$Vol_n(K + rB^n) = W_0(K) + \binom{n}{1}W_1(K)r$$
 (3)

$$+\binom{n}{2}W_2(K)r^2+\ldots+W_n(K)r^n.$$

The coefficient  $W_m(K)$  is called the m-th mean projection measure (Quermassintegral) of K.  $W_0(K) = \operatorname{Vol}_n(K)$ ,  $\binom{n}{1}W_1(K)$  is the surface area of K,  $W_n(K) = \operatorname{Vol}_n B^n = \kappa_n$ . In general, apart from a factor depending only on n,  $W_m(K)$  is the average of  $\operatorname{Vol}_{n-m}K(F)$  over all (n-m)-dimensional subspaces  $F \subseteq \mathbb{R}^n$ , where K(F) denotes the orthogonal projection of K on F. In particular,

$$W_{n-1}(K) = \frac{1}{2n} \int_{S^{n-1}} \operatorname{width}_n(K, x) \, dx, \tag{4}$$

<sup>\*</sup>Courant Institute, New York University, 251 Mercer Street, New York, NY 10012. This work was carried out while this author was at Rutgers University. Partially supported by NSF grant CCR-8902522.

<sup>&</sup>lt;sup>†</sup>Courant Institute, New York University, 251 Mercer Street, New York, NY 10012 and Mathematical Institute of the Hungarian Academy of Sciences, 1364 Budapest, Pf. 127. Research supported by NSF Grant CCR-89-01484, Hungarian Science Foundation Grant OTKA-1814 and the DIMACS Center.

where width<sub>n</sub> $(K, x) = Vol_1K(x)$  is the distance between the two supporting hyperplanes of K perpendicular to the

unit vector  $x \in S^{n-1}$ .

Let us turn now to the proof of Theorem 1. Imagine that the plane is embedded in  $\mathbb{R}^n$ . Put  $K = \text{conv}\{c_0, \ldots, c_n\}$ ,  $K' = \text{conv}\{c'_0, \ldots, c'_n\}$ . Furthermore, let  $B_i$  and  $B'_i$  denote the n-dimensional balls of radius rcentred at  $c_i$  and  $c_i'$  respectively. Obviously,  $\bigcup_{i=0}^n B_i \subseteq$  $K + rB^n$ .

Claim 1 Let diam K = $|c_i - c_j|$ . If r >max diam K, then

$$K + \left(r - \frac{(\operatorname{diam} K)^2}{r}\right) B^n \subseteq \bigcup_{i=0}^n B_i.$$

Proof: Let  $x \in K + (r - \frac{(\dim K)^2}{r})B^n$ , i.e, there exists  $k \in K$  such that  $|x - k| \le r - \frac{(\dim K)^2}{r}$ . If x is not a vertex of K, then one can choose  $c_i$  so that  $\angle xkc_i \le \frac{\pi}{2}$ .

$$|x-c_i| \le \sqrt{|x-k|^2 + |k-c_i|^2}$$

$$\le \sqrt{\left(r - \frac{(\operatorname{diam} K)^2}{r}\right)^2 + (\operatorname{diam} K)^2} \le r$$

provided that  $r \geq \operatorname{diam} K$ . Hence,  $x \in B_i$ .  $\square$ 

Claim 2 Let B<sub>i</sub> and B'<sub>i</sub> denote the n-dimensional balls of radius r centred at ci and c' respectively. Then,

$$\operatorname{Vol}_n(\bigcup_{i=0}^n B_i) \leq \operatorname{Vol}_n(\bigcup_{i=0}^n B_i').$$

Now we show how Theorem 1 can be deduced from these observations, and postpone the proof of Claim 2until the next section.
Combining Claims 1 and 2, we obtain that

$$\operatorname{Vol}_n\Big(K + (r - \frac{(\operatorname{diam} K)^2}{r})B^n\Big) \le \operatorname{Vol}_n(K' + rB^n).$$

Substituting (3), this implies

$$W_n(K)\Big(r-\frac{(\operatorname{diam} K)^2}{r}\Big)^n$$

$$+W_{n-1}(K)\left(r - \frac{(\operatorname{diam} K)^2}{r}\right)^{n-1} + O(r^{n-2})$$

$$\leq W_n(K')r^n + W_{n-1}(K')r^{n-1} + O(r^{n-2}).$$

Using the fact that  $W_n(K) = W_n(K') = \kappa_n$ , and taking the limits as r tends to infinity, we get

$$W_{n-1}(K) \le W_{n-1}(K'). \tag{5}$$

However, K is a planar convex set, so (4) can be rewritten as

$$W_{n-1}(K)$$

$$= \frac{1}{2n} \int_{S^1} \int_0^{+\frac{\pi}{2}} \int_{(\sin\phi)S^{n-3}} \operatorname{width}_2(K, y) \cos^2\phi \, dz d\phi dy$$

$$= \frac{(n-2)\kappa_{n-2}}{2n} \Big( \int_{S^1} \operatorname{width}_2(K, y) \, dy \Big)$$

$$\Big( \int_0^{+\frac{\pi}{2}} \sin^{n-3}\phi \cos^2\phi \, d\phi \Big).$$

Similarly,

$$W_{n-1}(K') = \frac{(n-2)\kappa_{n-2}}{2n} \left( \int_{S^1} \operatorname{width}_2(K', y) \, dy \right)$$
$$\left( \int_0^{+\frac{\pi}{2}} \sin^{n-3} \phi \cos^2 \phi \, d\phi \right).$$

Hence, it follows immediately from (5) that

$$\operatorname{Per} K = \frac{1}{2} \int_{S^1} \operatorname{width}_2(K, y) \, dy$$

$$\leq \frac{1}{2} \int_{S^1} \operatorname{width}_2(K', y) \, dy = \operatorname{Per} K',$$

completing the proof of Theorem 1.

# Proof of Claim 2.

The proof consists of two steps. The first one is a slight generalization of a simple fact which was also used by M. Gromov [G] to verify a special case of the following 'dual' counterpart of the Hadwiger-Kneser-Thue Poulsen conjecture: Let  $\{B_0, B_1, \ldots, B_m\}$ ,  $\{B_0', B_1', \ldots, B_m'\}$  be two sets of balls in *n*-dimensional space, and let  $c_i$  and  $c_i'$ denote the center of  $B_i$  and  $B'_i$ , respectively. If

$$|c_i - c_j| \le |c'_i - c'_j|$$
 for all  $i$  and  $j$ ,

then

$$\operatorname{Vol}_{n}(\bigcap_{i=0}^{m} B_{i}) \ge \operatorname{Vol}_{n}(\bigcap_{i=0}^{m} B'_{i}). \tag{6}$$

Gromov proved this result when the number of balls does not exceed n+1, i.e.,  $m \leq n$ .

**Lemma 1** Let  $K = \text{conv}\{c_0,\ldots,c_n\}$  and K' = $\operatorname{conv}\{c'_0,\ldots,c'_n\}$  be two non-degenerate simplices in ndimensional space, and assume that  $|c_i - c_j| \leq |c_i'|$  $c'_{i}$  for all i and j.

Then K can be continuously transformed into a congruent copy of K' in a finite number of steps so that

- 1. in each step we move only one vertex,
- 2. the motion of this vertex is smooth,
- 3. the edgelengths of the simplex never decrease.

Proof: Omitted in this abstract.

Lemma 2 Let  $\{B_0, B_1, \ldots, B_m\}$  be a collection of ndimensional balls of arbitrary radii in  $\mathbb{R}^n$ , and let  $c_i$  denote the center of  $B_i$ . Let  $\vec{\nu}$  be a vector with the property that translating  $c_0$  along  $\vec{\nu}$ , the distance between  $c_0$  and any other  $c_i$  does not decrease. Then  $\operatorname{Vol}_n(\bigcup_{i=0}^m B_i)$  does not decrease during this translation.

Proof: Omitted in this abstract.

Now we are in the position to prove Claim 2. Let us perturb slightly the point sets  $c_0, \ldots, c_n, c'_0, \ldots, c'_n \subseteq \mathbb{R}^n$  to make them full dimensional, without violating the con-

$$|c_i - c_j| \le |c'_i - c'_j|$$
 for all  $i$  and  $j$ .

Let us assume that this perturbation does not change  $\operatorname{Vol}_n(\bigcup_{i=0}^n B_i)$  and  $\operatorname{Vol}_n(\bigcup_{i=0}^n B_i')$  by more than  $\epsilon$ . Lemmata 1 and 2 now imply

$$\operatorname{Vol}_n(\bigcup_{i=0}^n B_i) - \epsilon \le \operatorname{Vol}_n(\bigcup_{i=0}^n B_i') + \epsilon$$

for any  $\epsilon > 0$ , and Claim 2 follows.

#### 3 Perimeter in the maximum norm.

Throughout we have been measuring distances in the usual Euclidean norm. However, it is possible that (2) is valid

for a large class of norms.

Let  $l_{\infty}$  denote the distance induced by the maximum norm, i.e., given two points  $a = (x, y), a' = (x', y') \in$  $\mathbf{R}^2$ ,  $l_{\infty}(a, a') = \max(|x - x'|, |y - y'|)$ . Let  $\operatorname{Per}_{\infty}$  denote the perimeter of a convex polygon measured in the  $l_{\infty}$ distance. We have the following

Theorem 2 Let  $\{c_0,\ldots,c_n\},\{c'_0,\ldots,c'_n\}$  be two point sets in the plane satisfying  $l_{\infty}(c_i, c_j) \leq l_{\infty}(c'_i, c'_j)$  for all i and i. Then

$$\operatorname{Per}_{\infty} \operatorname{conv}\{c_0,\ldots,c_n\} \leq \operatorname{Per}_{\infty} \operatorname{conv}\{c'_0,\ldots,c'_n\}.$$

Proof: First we make an observation about the perimeter of a convex polygon in the maximum norm.

Lemma 3 Given a finite point set P in the plane, let R denote the smallest rectangle enclosing P such that the angles between the sides of R and the coordinate axes are equal to  $\pi/4$ . Let a, b, c, d be (not necessarily distinct) elements of P such that each side of R contains at least one of them. Then

$$\operatorname{Per}_{\infty}\operatorname{conv}(P) = \operatorname{Per}_{\infty}\operatorname{conv}\left\{a,b,c,d\right\} = \frac{\operatorname{Per}(R)}{\sqrt{2}}.$$

Proof: Omitted in this abstract. Corollary. Let P be a finite point set in the plane,  $P^* \subseteq$ P. Then

$$\operatorname{Per}_{\infty} \operatorname{conv} P^* \leq \operatorname{Per}_{\infty} \operatorname{conv} P.\square$$

Next we show that it is sufficient to prove Theorem 2 for n = 3 (for 4 points). For n < 3 there is nothing to prove. Assume that the statement is true for n = 3, and sitting on the boundary of the rectangle R enclosing P, as in Lemma 3. Let a', b', c', d' denote the corresponding points in  $P' = \{c'_0, \ldots, c'_n\}$ . Suppose, in order to obtain a contradiction, that

 $\operatorname{Per}_{\infty} \operatorname{conv} P > \operatorname{Per}_{\infty} \operatorname{conv} P'$ .

By Lemma 3 and by the above Corollary.

$$\operatorname{Per}_{\infty} \operatorname{conv} \{a, b, c, d\} = \operatorname{Per}_{\infty} \operatorname{conv} P >$$

$$\operatorname{Per}_{\infty} \operatorname{conv} P' \geq \operatorname{Per}_{\infty} \operatorname{conv} \{a', b', c', d'\},$$

contradicting our assumption that Theorem 2 is true for

n = 4. Suppose now that in the maximum norm distances between the elements of  $P = \{a, b, c, d\}$  are at most as large as the corresponding distances within  $P' = \{a', b', c', d'\}$ .

$$Per_{\infty} conv P > Per_{\infty} conv P'$$
.

We can assume without loss of generality that a', b', c', d' are in convex position. Suppose not. Then one of them, say d', is in the interior of the triangle induced by the others. The  $l_{\infty}$ -distance of d' from a', b' and c' is either their vertical distance or their horizontal distance. Suppose without loss of generality that at most one of these three distances is the horizontal distance. Then we can move d' horizontally until it hits the boundary of the

triangle a'b'c', so that none of the  $l_{\infty}$ -distances decreases. We can also assume that a, b, c, d are in convex position. Otherwise, if, say, d is in the triangle abc, then by the

Corollary

$$Per_{\infty} conv \{a, b, c\} = Per_{\infty} conv P$$

$$> \operatorname{Per}_{\infty} \operatorname{conv} P' \ge \operatorname{Per}_{\infty} \operatorname{conv} \{a', b', c'\},$$

a contradiction. Let abcd be the cyclic order of the vertices of convP. By symmetry, there are only two essentially different cases.

• Case A: The cyclic order of the vertices of conv P' is a'b'c'd'. Then

$$\operatorname{Per}_{\infty} \operatorname{conv} P = l_{\infty}(a,b) + l_{\infty}(b,c) + l_{\infty}(c,d) + l_{\infty}(d,a)$$

$$\leq l_{\infty}(a',b') + l_{\infty}(b',c') + l_{\infty}(c',d') + l_{\infty}(d',a')$$
  
=  $\operatorname{Per}_{\infty} \operatorname{conv} P'$ .

• Case B: The cyclic order of the vertices of conv P' is a'c'b'd'. By the triangle inequality for the maximum norm,  $l_{\infty}(a,b)+l_{\infty}(c,d) \leq l_{\infty}(a,c)+l_{\infty}(b,d)$ . Hence,

Per<sub>\infty</sub> conv 
$$P = l_{\infty}(a,b) + l_{\infty}(b,c) + l_{\infty}(c,d) + l_{\infty}(d,a)$$

$$\leq l_{\infty}(b',c') + l_{\infty}(d',a') + l_{\infty}(a',c') + l_{\infty}(b',d')$$

$$= \operatorname{Per}_{\infty} \operatorname{conv} P'.$$

This completes the proof of Theorem 2.

# Related problems and generalizations.

Theorem 1 can readily be generalized to bounded infinite sets  $S \subset \mathbb{R}^2$ . A mapping  $f: S \to \mathbb{R}^2$  is said to be a

$$|f(p)-f(q)| \le |p-q|$$
 for all  $p, q \in S$ .

Theorem 3 Let f be a contraction of a bounded set  $S \subseteq$ R<sup>2</sup>. Then

$$\operatorname{Per conv} f(S) \leq \operatorname{Per conv} S.\square$$

Evidently, similar results cannot be true for the surface area of the convex hull of higher dimensional sets. However, the above arguments immediately yield

Theorem 4 Let f be a contraction of a bounded set  $S \subseteq$  $\mathbb{R}^n$ . Then

$$W_{n-1}(\operatorname{conv} f(S)) \le W_{n-1}(\operatorname{conv} S),$$

where  $W_{n-1}$  is the (n-1)-st mean projection measure (the mean width) in n dimensions.

It might be interesting to notice that by slightly modifying the proof presented in Section 2, we can generalize Lemma 2 in two different directions.

Lemma 2' Let  $\{B_1, \ldots, B_k, B_{k+1}, \ldots, B_m\}$ , k < m be a collection of n-dimensional balls of arbitrary radii in  $\mathbb{R}^n$ , and let  $c_i$  denote the center of  $B_i$ . Let  $\vec{v}$  be a vector with the property that translating  $\{c_1,\ldots,c_k\}$  along  $\vec{v}$ , the distance between any  $c_i$  and  $c_j$ ,  $i \le k < j$  does not decrease. Then

(i) 
$$\operatorname{Vol}_n(\bigcup_{i=1}^m B_i)$$
 does not decrease,

(i) 
$$\operatorname{Vol}_n(\bigcup_{i=1}^m B_i)$$
 does not decrease,  
(ii)  $\operatorname{Vol}_n(\bigcap_{i=1}^m B_i)$  does not increase,

during this translation.

Combining Lemma 1 and (the special case k = 1 of) Lemma 2'(ii), we obtain a proof of (6) for  $m \leq n$ , somewhat different from the one given in [G].

It is worth mentioning that conjecture (1) of Thue Poulsen, Kneser and Hadwiger cannot be generalized to other Minkowski planes. More precisely, we have the fol-

Theorem 5 Let the plane be equipped with a norm ||.|| such that the unit disk  $C = \{x \in \mathbb{R}^2 : ||x|| \le 1\}$  is not

Then one can find points  $a, b, a', b' \in \mathbb{R}^2$  with the property that ||a - b|| < ||a' - b'|| but

$$Area((C+a) \cup (C+b)) > Area((C+a') \cup (C+b')).\Box$$

On the other hand, the weaker statement (2) might remain true for a large class of other norms substantailly different from the Euclidean one.

In the last section we have shown that (2) holds in the plane equipped with the maximum norm (when the unit disk is a square). A related question is the following.

Given two collections of axis parallel unit squares in the plane,  $S_1, \ldots, S_n$  and  $S'_1, \ldots, S'_n$ , such that  $Area(S_i \cup S_j) > Area(S'_i \cup S'_j)$  for all i and j. Is it true that  $Area(\cup S_j) > Area(\cup S'_j)$ ?

Some related questions with applications to particle physics are discussed by Lieb and Simon in [L] and [LS].

#### References

- [A] R. Alexander, Lipschitzian mappings and total mean curvature of polyhedral surfaces I, Trans. of American math. soc. 288 (1985) No. 2, 661-678.
- T. Bonnesen and W. Fenchel, Theorie der Kon-[BF] vexen Körper, Springer, Berlin, 1934 (reprinted by Chelsea Publ. Co., New York, 1948).
- B. Bollobás, Area of the union of disks, Elem. d. Math. 23(1968), 60-61.
- [Bu] H. Busemann, Convex Surfaces, Interscience Publ.. London-New York, 1958.
- [G] M. Gromov, Monotonicity of the volume of intersection of balls. In: Geometrical Aspects of Functional Analysis, Lecture notes in Mathematics, Springer-Verlag, Berlin 1267(1987), 1-4.
- [H] H. Hadwiger, Ungelöste Probleme No. 11, Elem. d. Math. 11(1956), 60-61
- [K] M. Kneser, Einige Bemerkungen über das Minkowskische Flächenmass, Arch. Math. 6(1955), 382-390.
- [L]K. Leichtweiss, Konvexe Mengen, Springer Verlag, Berlin-Heidelberg-New York, 1980.
- [Li] E. Lieb, Monotonicity of the molecular electronic energy in the nuclear coordinates J. Phys. B: At. Mol. Phys. 15(1982), L63-L66.
- E. Lieb and B. Simon, Monotonicity of the elec-[LS] tronic contribution to the Born-Oppenheimer energy J. Phys. B: At. Mol. Phys. 11(1978), L537-L542.
- [S] V. Sudakov, Gaussian Random processes and measures of solid angles in Hilbert space Soviet Math. Dokl. Vol. 12(1971) No.2
- [TP] E. Thue Poulsen, Problem 10, Math. Scand. 2(1954), 346.