# Shortest Watchman Tours in Weak Visibility Polygons

P.Pradeep Kumar\*and C.E.Veni Madhavan<sup>†</sup>
Department of Computer Science and Automation
Indian Institute of Science, Bangalore, India

June 17, 1993

#### Abstract

An  $O(n^4 log log n)$  algorithm for shortest watchman tour (SWT) problem for simple polygons, given a starting point on the boundary of the polygon is proposed in Chin and Ntafos [CN1]. The problem of finding the SWT in general polygons when the starting point is not specified is open. We develop an  $O(n^{10})$  algorithm for the SWT problem in weak visibility polygons, with no assumption on the starting point.

#### 1 Introduction

The watchman tour problem was introduced by Chin and Ntafos [CN2]. Earlier, many researchers considered the problem of positioning stationary watchmen in a gallery so that every point in the gallery can be seen by atleast one watchman. The problem of finding the minimum number of watchmen needed is equivalent to that of covering the polygon with minimum number of star-shaped components.

Chin and Ntafos [CN2] considered the problem of finding a watchman tour, which is a tour within a polygon with the property that every point in the polygon is visible from at least one point along the tour. They present O(n) and O(nlogn) algorithms to find a shortest watchman route in a simple rectilinear monotone and simple rectilinear polygon, respectively. Here n is the number of vertices in the polygon. Chin and Ntafos [CN1] developed an  $O(n^4loglogn)$  algorithm to find a shortest watchman route in a simple polygon given a starting point on its boundary. An interesting incremental algorithm for the SWT problem gas recently been proposed [HIT].

We present a polynomial time algorithm for finding the shortest watchman tour, with no assumption on the starting point, for a class of polygons called weak visibility polygons.

Consider a simple polygon P. Two points in P are said to be visible if the line segment joining them lies totally inside P. A point p is said to be weakly visible from an edge st if there is a point z in the interior of st such that p and z are visible. If every point in P is weakly visible from st then P is called a weak visibility polygon. If a polygon P is weakly visible from an edge, then that edge is called a weak visibility edge. Pal [SP90] presents characterizations of weak visibility polygons, and based on them, an algorithm for recognizing weak visibility polygons with running time proportional to the size of the

<sup>\*</sup>presently at CS dept. Ohio State Univ., Columbus, USA

presently visiting DIMACS Center, Rutgers University, USA, cevm@dimacs.rutgers.edu

visibility graph of the input polygon. Here, visibility graph is the graph with the vertices of the polygon as the vertex set, and two vertices are adjacent if they are visible from each other in the polygon.

Assume that the simple polygon P is given as a clockwise sequence of vertices  $v_1, v_2, ... v_n$  with their x and y coordinates.  $\overline{v_i v_{i+1}}$  are the edges. The direction of edge  $\overline{v_i v_{i+1}}$  is assumed to be from  $v_i$  to  $v_{i+1}$ .  $v_i$  is said to be the predecessor of  $v_{i+1}$ , denoted by  $\operatorname{pred}(v_{i+1})$  and  $v_{i+1}$  is said to be the successor of  $v_i$  denoted by  $\operatorname{succ}(v_i)$ . It is assumed that no three consecutive vertices are collinear. Denote the boundary of P by  $\operatorname{bd}(P)$ , a weak visibility edge by  $\operatorname{WV}E$  and the shortest watchman tour by  $\operatorname{SWT}$ . Assume that a  $\operatorname{WV}E$  of P is given and that P is aligned with the  $\operatorname{WV}E$  along the x-axis.

# 2 Properties of SWT and related results.

In what follows we develop through a sequence of definitions and propositions, the properties satisfied by candidate tours which include the SWT. These propositions characterize the corners or turning points and the segments of such tours. We suppress the proofs in this version.

Visibility polygon of a vertex v is the portion of the given polygon P, denoted VP(v), such that v is visible to all points in VP(v). The boundary of VP(v) consists of segments through the interior of P, alternating with chains of boundary segments of P. The segments through the interior of P, if extended through the interior of the polygon, will intersect at v. VP(v) has at least one point on the WVE, since every vertex v is visible from the WVE. Adjacent segments of a vertex are the two segments on the bd(P) meeting at the vertex. A chord is a segment connecting two points on bd(P) through the interior of P. Supporting Chords of a vertex v, are the extensions of the adjacent segments of v, away from it through the interior of P until they meet the boundary of the polygon. In some cases, one or more supporting chords may not exist, in which case, they are assumed to have zero length. Supporting Chains of a vertex v are the bounding chains of the visibility polygon of v, starting from  $v_i$  and  $v_j$  where  $v_iv$  and  $vv_j$  are the adjacent segments of v. The supporting chains can contain the boundary edges of the polygon P. They necessarily include the supporting chords of v.

**Proposition 2.1** A supporting chain of any vertex v can intersect a segment  $\overline{v_1v_2}$  lying entirely within the interior of P, in atmost one point.

Directions of edges and segments of supporting chains: To each edge on bd(P), we can assign a direction as described before. Each supporting chord is assigned a direction identical to that of the adjacent segment, which when extended, gives the supporting chord. Each segment in a supporting chain C has direction identical to that of the adjacent segment from which C starts.

Inner region of a chord is the region in P to the right, while travelling along its assigned direction. Analogously, the outer region is the region to the left. Inner region of a chain of segments is the intersection of the inner regions of all the segments of the chain. Outer region of a chain is the region in P other than the inner region. It can be seen that the visibility polygon of a vertex is the intersection of the inner regions of supporting chains of the vertex and the boundary of the polygon. A chord s of P is said to cover a chord t if t lies entirely in the outer region of s. r-supporting chord of a vertex v above the WVE is that supporting chord of v whose direction is towards v and t supporting chords are those supporting chords whose directions are away from v and towards v, respectively. Two supporting chords are said to be of the same kind if both are t-supporting or t-supporting. Two segments are said

to be of the same kind if they are parts of supporting chords of the same kind. They are said to be of the *opposite* kind, otherwise. The set of l-extreme chords is the subset of l-supporting chords which are not covered by any other l-supporting chord. Similarly, the set of r-extreme chords is the subset of r-supporting chords which are not covered by any other r-supporting chord.

Proposition 2.2 In a weak visibility polygon, the cover relation is asymmetric and transitive for supporting chords of the same kind. It means that if an l- supporting chord s covers an l- supporting chord t, t cannot cover s, and hence, s should be in the inner region of t. Also, if s covers t and t covers v, then s should cover v. Similar results hold for r- supporting chords.

If the outer regions of two extreme chords of the same kind intersect, then either one of them covers the other or the two chords intersect.

**Proposition 2.3** The entire polygon P is visible from a continuous tour in P iff all the convex vertices of P are visible from the tour.

The extreme chords cover all other chords of the supporting chains of every convex vertex.

All the convex vertices and hence the whole polygon is visible to a watchman tour visiting the visibility polygons of vertices whose supporting chains contain at least one extreme chord.

A corner of a tour is defined to be a point on the tour at which the tour changes direction. We label the corners of a tour in increasing order in clockwise direction. A corner of a tour is said to be convex if the angle swept by rotating the segment  $vv_i$  to the segment  $vv_j$ , about v in a counterclockwise direction is less than 180 degrees. Here  $v_i$  and  $v_j (i < j)$  are the corners preceding and succeeding v in the tour. It is said to be reflex, otherwise. A tour is said to reflect at a point p on a supporting chord s if p is on the chord and both corners of the tour adjacent to p are in the outer region of s. The segments of the l-extreme and r-extreme chords on which a SWT can reflect are termed l-reflecting, and r-reflecting segments, respectively.

Proposition 2.4 Each convex corner of the SWT is a point of reflection on a supporting chain or a point on the boundary of the polygon, and any reflex corner of the tour is a point on the boundary of the polygon.

**Proposition 2.5** A SWT reflecting on a supporting chord should lie entirely in the outer region of the chord. It cannot touch or intersect the chord at any point other than the point of reflection. (The SWT can graze along a supporting chord).

A SWT can reflect only on extreme chords.

The SWT lies entirely in the intersection R of outer regions of all extreme chords on which it reflects and the reflecting segments are on the boundary of this region.

An essential chord is an extreme chord whose outer region contains portions of each of the visibility polygons having one or more extreme chords on their boundaries. Call the edges on the boundary of R containing the reflecting segments as essential segments.

**Proposition 2.6** All points of reflection of the SWT are on essential chords. All essential chords of the same kind, as an essential chord s on which the SWT reflects, should intersect s.

**Proposition 2.7** The l-essential and r-essential segments (containing l-reflecting, and r-reflecting segments, respectively) form two distinct convex chains on the boundary of R.

The corners of the SWT which are points of reflection on l-reflecting and r-reflecting segments form two continuous convex chains.

# 3 The Algorithm

### 3.1 Finding essential segments

The directions of the supporting chords can be assigned by a clockwise scan around the boundary of the polygon. The extreme chords can be determined by checking pairs of supporting chords and eliminating one of them if it is covered by the other. Essential chords can be found by checking for each extreme chord whether its outer region contains portions of visibility polygons corresponding to all other extreme chords. This can be done by checking whether the extreme chord is in the inner region of or intersects both the supporting chains of the visibility polygons corresponding to the extreme chords. If so, then it means that the outer region of the extreme chord contains some portion of the visibility polygon and that the extreme chord is essential.

### 3.2 Finding Reflecting Segments

Let L and R be the sets of l-essential and r-essential chords. We traverse the SWT in an anticlockwise direction. Let  $f_l$ ,  $e_l$  denote the first and last l-extreme chords on which the SWT reflects, and let  $f_r$  and  $e_r$  be similarly the first and last r-extreme chords on which the SWT reflects.

#### 3.2.1 Selecting $f_l$ and $e_l$

Due to intersections with other supporting chords, the chords in L are divided into  $O(n^2)$  segments (each supporting chain can intersect a chord in at most one point, by Proposition 2.1). It is required to select the  $f_l$  and  $e_l$  from these.

Consider the intersections among the l-extreme chords alone. Each l-extreme chord can intersect all the others. We traverse along these chords in their assigned direction. The segments of all the chords until the first intersection are candidates for  $f_l$ . Subsequently, at each intersection, all except one of the intersecting extreme chords can be ignored (i.e., the rest of the segments in those chords cannot be candidates for  $f_l$ ), since a tour reflecting first on any segment on these chords will fail to visit the visibility polygon corresponding to at least one of the extreme chords intersecting at the point. Thus, there can be a maximum of 2m+1 segments remaining where m is the number of chords in L, where  $m \le n$ . Due to intersections with other supporting chords, each of these segments may be split into O(n) segments (each supporting chain can intersect the segment in at most one point). Each of these is a candidate for  $f_l$  and hence, there are  $O(n^2)$  candidates for  $f_l$  where n is the number of vertices of the polygon.

Similarly, the first segments while travelling along the l-extreme chords against their assigned direction are candidates for  $e_l$ . At each intersection, the subsequent segments of all except one l-extreme chords intersecting at that point can be ignored since, if the SWT reflects last on one of these segments, the visibility polygons corresponding to the other l-extreme chords at the intersection cannot be visited. There are  $O(n^2)$  candidates for  $e_l$  by an argument similar to that for l-extreme chords.

The case of  $f_r$  and  $e_r$  is similar.

# 3.3 Finding Reflecting Chains

The SWT reflects either on r-essential chords alone, or on l-essential chords alone, or on both r and l essential chords, depending on the input polygon. Let  $F_l$ ,  $F_r$  and  $E_l$ ,  $E_r$  be the extreme chords containing

 $f_l$ ,  $f_r$  and  $e_l$ ,  $e_r$  respectively. If the set  $\{f_l, e_l, f_r, e_r\}$  corresponds to a shortest watchman tour, it should be the case that the outer regions of  $F_l$ ,  $E_l$ ,  $F_r$  and  $E_r$  intersect in a non-empty region, since the SWT lies in the outer region of all the extreme chords on which it reflects. Since the outer regions of  $F_l$  and  $E_l$  intersect,  $F_l$  should intersect  $E_l$  since none of them covers the other, by Proposition 2.3. Similar is the case with  $F_r$  and  $E_r$ . The SWT has to reflect on or intersect all extreme chords l intersecting  $F_l$  and  $E_l$  such that  $f_l$  and  $e_l$  are in the outer region of l, and l passes through the intersection of the outer regions of  $F_l$  and  $E_l$ , since otherwise, the watchman tour cannot visit the visibility polygon corresponding to l. Call these chords the candidate reflecting chords for  $(f_l, e_l)$ . Similar chords in the case of  $F_r$  and  $E_r$  also have to be intersected or reflected upon by the watchman tour. Define the included angle between x and y to be the angle formed by rotating a line segment from x to y in an anti-clockwise direction denoted Includedangle(x, y). Let  $\theta$  be the included angle between  $F_l$  and  $E_l$ . Let  $C = \langle l_1, l_2, l_3, \ldots \rangle$  be the ordered set of reflecting chords, for  $(f_l, e_l)$  such that  $x_1, x_2, \ldots, x_n$ , their corresponding intersections with  $F_l$  are in the order in which they appear while travelling along  $F_l$  in its assigned direction. Let  $\alpha_1, \alpha_2, \ldots, \alpha_n$  be the included angles made by the candidate reflecting chords  $l_1, l_2, \ldots, l_n$  respectively with  $F_l$ .

**Proposition 3.1** (a) The SWT should reflect on  $R = \langle y_1, y_2, ..., y_k \rangle$  the maximum sized subset of  $\langle l_1, l_2, ..., l_n \rangle$  such that:

- 1.  $\langle y_1, y_2, \ldots, y_k \rangle$  appears in the same relative order in R as in C,
- 2.  $Included angle(F_l, y_l) < \dots Included angle(F_l, y_k) < Included angle(F_l, E_l)$ .
- 3. There does not exist a chord  $y_m \in \mathbb{C}$  and  $y_m \notin \mathbb{R}$  such that  $y_n < y_m$  and Includedangle $(F_l, y_m) \leq Includedangle(F_l, y_n)$  for any  $y_n \in \mathbb{R}$ .

In other words, for any  $y_j \in \mathbb{R}$ , there does not exist  $y_k \in \mathbb{R}$ , k < j and Includedangle $(F_l, y_k) > Includedangle(F_l, y_j)$ .

- (b) The set  $\mathbf{R}$  is unique and can be constructed by moving down  $F_l$  along its direction and, when an intersection  $\mathbf{z}_i$  with an  $l_i \in \mathbf{C}$  is encountered, removing all candidate reflecting chords  $l_k$  intersecting  $F_l$  before  $\mathbf{z}_i$  such that Includedangle( $F_l, y_k$ ) > Includedangle( $F_l, y_i$ ).
  - (c) All chords in C, but not in R are intersected by any path reflecting on the chords of R.

# 3.4 Checking Validity of Reflecting Chains

A set of reflecting segments is valid if there exists a tour reflecting on these segments and visiting the visibility polygons corresponding to all the extreme chords.

# Proposition 3.2 A set S of reflecting segments is valid

only if each of the reflecting segment is in the outer region of the extreme chord corresponding to each of the other reflecting segments, and, if so,

if and only if for each vertex v such that atleast one of its supporting chords is an extreme segment containing no reflecting segment, there exist atleast one of the reflecting segments of s to the inner regions of each of the supporting chains of v.

### 3.5 Finding the shortest length tour reflecting on prescribed segments

The reflecting chain(s), along with the boundary of the polygon form a polygon which consists of a chain of one or more simple polygons, with each simple polygon connected to the next by a single point. The reflecting segments are on the boundary of the terminal polygons. If the chain has more than one simple polygon, the SWT has to pass through the points connecting the polygons, and it is sufficient to find the minimum length tour reflecting on the prescribed segments of the terminal polygons, passing through the interconnecting points to the next polygon in the chain, and finally connect the tours of the terminal polygons by the shortest tour through the interconnecting points of the chain. We use an O(nloglogn) algorithm of [CN2] which first triangulates the polygon, and considering the reflecting segments as 'mirrors', 'unfolds' the polygon and finds the shortest path connecting one of the reflecting segments to its 'reflected image'.

#### 4 Proof of Correctness

**Proposition 4.1** The algorithm - shortest watchman tour computes in polynomial time, the SWT in a given weak visibility polygon.

#### Proof

The reflecting segments form one or two continuous, convex chains according to Proposition 2.7. The method in Section 3.2 selects all the possible first and last segments for these chains. Proposition 3.1 proves the method for finding the unique convex chains between the selected corner segments of the same type. By Proposition 2.3, it is sufficient for the watchman tour to visit the visibility polygons corresponding to the extreme chords. Hence, by Proposition 3.2 the algorithm computes all possible watchman tours for the polygon, and hence, the Shortest Watchman Tour.

Let n be the number of convex vertices. Finding extreme chords can be done in  $O(n^2)$  time (each pair of supporting chords may have to be considered to check whether one of them is covered by the other and there are  $O(n^2)$  of them). Since there are  $O(n^2)$  candidates for  $f_l, e_l, f_r$  and  $e_r$ , there are  $O(n^4)$  possible  $(f_l, e_l)$  and  $(f_r, e_r)$  pairs, and  $O(n^8)$  quadruples  $(f_l, e_l, f_r, e_r)$ . Checking for valid chains take  $O(n^2)$  time and finding the minimum length touching the prescribed reflecting segments take  $O(n\log\log n)$  time. Thus, the overall complexity of the algorithm is  $O(n^{10})$ .

## References

- [1] [SP] Sudebkumar Prasant Pal, 'Weak Visibility and related problems on simple polygons', Ph.D. thesis, 1990, Indian Institute of Science, Bangalore, India.
- [2] [CN1] W.P.Chin and S.Ntafos, 'Shortest Watchman Routes in simple polygons', Tech. Report.
- [3] [CN2] W.P.Chin and S.Ntafos, 'Optimum Watchman routes', Info. Proc. Lett. 28, 1988, 39-44.
- [4] [HIT] Hirata, Inagaki, and Tan, 'An incremental algorithm for constructing shortest watchman tours' Int. Symp. on Algorithms ISA 91, LNCS 557, 1991.