Tamal K. Dey<sup>1</sup>

#### Abstract

Given a cycle of length k on a triangulated 2-manifold, we determine if it is null-homologous (bounds a surface) in O(n+k) optimal time and space where n is the size of the triangulation. Further, with a preprocessing step of O(n) time which is performed only once, we answer the same query for any cycle of length k in O(g+k) time. This is optimal for k>g.

### 1 Introduction

In recent years a new focus has developed in studying the algorithmic aspects of topology [2, 5, 6, 7], a well developed branch of mathematics. This emergent field has been called "Computational Topology" [5, 7]. It is generally recognized that there is a vast repository of topological problems which have not been studied extensively from algorithmic point of view. In this paper, we address one such problem, namely, topologically distinguishing the curves(cycles) on 2-manifolds that bounds a surface (possibly empty and is not necessarily a disk). These cycles are called *null-homologous* cycles.

The importance of detecting null-homologous cycles comes from two facts. Homology groups of a topological space reveals its connectivity. To compute them efficiently Delfinado and Edelsbrunner [2] showed a geometric approach. Unfortunately this approach did not have any efficient implementation in dimensions higher than three since there is no known polynomial time algorithm to detect nullhomologous cycles in higher dimensions. Although this paper does not provide a solution for this general problem, it gives a better understanding of null-homologous cycles on manifolds. Secondly, dectecting null-homologous cycles is related to a more difficult problem called contarctability problem, which asks if a given cycle is contractable to a single point. These cycles are called null-homotopic cycles. For 2-manifolds all null-homotopic cycles are null-homologous, though the reverse is not true. Recently, we provided an improved algorithm for detecting null-homotopic cycles on 2-manifolds. This algorithm runs in O(n+gk) time where n is the size of the triangulation, k is the cycle length, and g is the genus of the given 2-manifold. We can use the algorithm for null-homologous cycles first, and declare those cycles not null-homotopic that are not null-homologous. Of course, we have to run the algorithm for null-homotopic cycles when a cycle is detected to be null-homologous. This approach saves time in those cases where the cycle is not null-homologous since null-homologous cycles can be detected in O(n+k) time as shown here.

## 2 Preliminaries

A 2-manifold is a topological space in which each of its points has a neighborhood homeomorphic to an open disk. A 2-manifold can be infinite or finite. Moreover, it can be closed or open depending

<sup>&</sup>lt;sup>1</sup>Computer Science, Indiana University-Purdue University at Indianapolis, Indianapolis, IN 46202, USA.

on whether its closure coincides with itself or not. A closed and bounded 2-manifold is also called a compact 2-manifold. A Sphere and a klein bottle are two examples of compact 2-manifolds. A 2-manifold is called orientable if it has two distinct sides. Otherwise, it is non-orientable. For details see [4]. In this paper we consider only compact 2-manifolds.

A 2-manifold is triangulable in the sense that it can be represented as the union of a set of triangles, edges and vertices satisfying the following properties. Each pair of triangles either share a single vertex or a single edge, or are completely disjoint. Also, the triangles incident on a vertex can be ordered circularly so that two triangles share a common edge if and only if they are adjacent in this ordering.

#### 2.1 Chains and Boundaries

A precise definition of null-homologous cycles can be given algebraically. We use the notations of [4] for this purpose. Although we will be working with the triangulations of 2-manifolds which are simplicial 2-complexes, we define all terms here in the most general setting. Let K be a simplicial complex. An oriented simplex  $[v_0, v_1, ..., v_p]$  is the p-simplex with vertices  $v_0, v_1, ..., v_p$  and the particular ordering on these vertices,  $v_0v_1...v_p$ . A p-chain on K is a function c from the set of oriented p-simplices of K to the integers where

- 1.  $c(\sigma) = -c(\sigma')$  if  $\sigma$  and  $\sigma'$  are opposite orientations of the same simplex.
- 2.  $c(\sigma) = 0$  for all but finitely many oriented p-simplices  $\sigma$ .

For an oriented simplex  $\sigma$ , the elementary chain c is the function defined as

```
c(\sigma) = 1

c(\sigma') = -1 if \sigma' is the opposite orientation of \sigma.

c(\tau) = 0 for all other oriented simplices \tau.
```

We use the symbol  $\sigma$  to denote not only a simplex, or an oriented simplex, but also to denote the elementary p-chain c for the oriented simplex  $\sigma$ . With this notation we write a p-chain d as  $d = a_1\sigma_1 + a_2\sigma_2 + ... + a_k\sigma_k$  where  $a_i$ 's are integer coefficients.

If  $\sigma = [v_0, v_1, ..., v_p]$  is an oriented simplex with p > 0, we define  $\delta_p \sigma = \sum_{i=0}^p (-1)^i [v_0, ..., \hat{v_i}, ..., v_p]$ , where the symbol  $\hat{v_i}$  means that the vertex  $v_i$  is to be deleted from the array. The interpretation of  $\delta_p$  is that it finds the oriented boundary of an oriented simplex. For example, an oriented triangle  $[v_0v_1v_2]$  gives the oriented cycle  $[v_0v_1]+[v_1v_2]+[v_2v_1]$  when operated by  $\delta_2$ . For a p-chain  $d=\sum a_i\sigma_i$ ,  $z=\delta_p d=\sum a_i\delta_p\sigma_i$  is called the (p-1)-boundary of d. We also say z bounds d. A p-chain d is called a p-cycle if  $\delta_p d=0$ .

With the addition p-chains form a group  $C_p(K)$ , called the group of p-chains of K. If p < 0 or p > dim K, we let  $C_p(K)$  denote the trivial group. These groups are abelian [4]. The function  $\delta_p: C_p(K) \to C_{p-1}(K)$  is an homomorphism. The set of p-chains that are mapped by  $\delta_p$  to the identity element of  $C_{p-1}(K)$  are called the group of p-cycles and denoted  $Z_p(K)$ . In other words,

the kernel of  $\delta_p$  is  $Z_p(K)$ . The image of  $\delta_{p+1}:C_{p+1}(K)\to C_p(K)$  is called the group of p-boundaries and is denoted  $B_p(K)$ . A careful observation reveals that  $B_p(K)\subset Z_p(K)$ . The quotient group  $H_p(K)=Z_p(K)/B_p(K)$  is called the pth homology group of K. A p-cycle z in K is null-homologous if it represents an identity in  $H_p(K)$ . Stated otherwise, a p-cycle z is null-homologous if there exists a p+1-chain d so that z bounds d. So a 1-cycle z on a triangulated 2-manifold is null-homologous if there is a 2-chain d bounded by z. Two p-cycles  $z_1, z_2$  are called homologous if there exists a p+1-chain d so that  $z_1-z_2=\delta_p d$ .

## 2.2 Polygonal Schema

Any orientable or non-orientable 2-manifold can be represented by a simple polygon P with even number of edges which is also called a polygonal schema of S. Each edge of P has a signed label such that each unsigned label occurs twice. See [3] for details. Two edges with the same unsigned labels are called partnered edges. Partnered edges can have labels with the same or opposite signs. Two partnered edges with labels +x and -x represent the same edge on S but are oppositely directed on P. The labels of the edges that are directed in a clockwise direction around P are signed positively. In general, we use  $x^c$  to denote the complement of the label x. To reconstruct a surface homeomorphic to S from this polygonal representation, the oriented edges with the same labels are identified together in such a way that their orientations match. For simplicity, we say that S is obtained from P by identifying partnered edges appropriately.

An orientable 2-manifold with genus g > 0 can be represented canonically by a 4g-gon where the labels on the edges around the polygon are of the form:  $x_1y_1x_1^cy_1^cx_2y_2x_2^cy_2^c...x_gy_gx_g^cy_g^c$ . For g = 0, the 2-manifold is a sphere which can be represented canonically by two directed edges  $xx^c$ .

Similarly, a non-orientable 2-manifold can be represented canonically by a 2g-gon where the labels on the edges around the polygon are of the form:  $x_1x_1x_2x_2...x_gx_g$ .

# 2.3 Cycles and Null-homology

Let T denote a triangulation of an orientable 2-manifold S. Let C be any given cycle (oriented) on T. The cycle C is a sequence of oriented edges with the first and the last edge meeting at a vertex. Let P be a polygonal schema of S with a triangulation T' such that there is a one-to-one correspondence between triangles of T and T'. The following lemma proves that such a polygonal schema can be constructed.

LEMMA 2.1 A polygonal schema P with triangulation T' can be constructed from T where there is a one-to-one correspondence between triangles of T' and T.

PROOF. We construct a sequence of closed disks  $D_1, D_2, ..., D_n$  in a plane incrementally such that  $P = D_n$  at the end. Initially,  $D_1 = \sigma'$ , a triangle in the plane that corresponds to an arbitrarily chosen triangle  $\sigma_1$  on M. Let  $D_i = \sigma'_1 \cup \sigma'_2 ... \cup \sigma'_i$  after the *ith* step. At the i+1 th step we choose a triangle  $\sigma_{i+1}$  on T which has the following properties: (i) no triangle corresponding to  $\sigma_{i+1}$  has been included in  $D_i$ , and (ii) a triangle  $\sigma_j$  adjacent to  $\sigma_{i+1}$  by an edge has a corresponding triangle

in  $D_i$ . These two conditions imply that there is an edge  $e = \sigma_j \cap \sigma_{i+1}$  such that its corresponding edge e' on  $D_i$  appears on  $bd(D_i)^2$ . We attach a triangle  $\sigma'_{i+1}$  to  $bd(D_i)$  such that  $\sigma'_{i+1} \cap bd(D_i) = e'$ . This gives the new disk  $D_{i+1} = D_i \cup \sigma'_{i+1}$ . It is clear that if  $D_i$  is a closed disk, so is  $D_{i+1}$ . Finally, when we exhaust all triangles on M, we have  $D_n$  with the desired triangulation T'.

Let G be the graph on T obtained by identifying the partnered edges of bd(P). If C intersects the edges of G, C can be written as a sequence of directed paths in T' with end points on bd(P). Let  $U = \{u_1, u_2, ..., u_r\}$  be this sequence of paths. A directed path  $u_i$  from  $v_1$  to  $v_2$  can be deformed to another directed path  $u_i'$  that lies entirely on bd(P) and to the left of  $u_i$ . Let  $U' = \{u_1', u_2', ..., u_r'\}$ . The paths in U' correspond to a cycle C' on G. One can think of this process as deforming C to another cycle C' on T so that all edges of C' lie on the edges of G. We also say that C' is carried by bd(P). In terms of homology, C and C' are homologous since C - C' is a 2-chain consisting of the triangles between  $u_i$  and  $u_i'$  for i = 1, ..., r. If C does not intersect G, then it is trivially null-homologous since the corresponding cycle in T' lies in  $int(P)^3$  which is an open disk. Henceforth, we consider only the difficult case where C intersects G.

Let  $x_1, x_2, ..., x_t$  be the edges on G. Let  $a_1, a_2, ..., a_{2t}$  be the sequence of signed labelled edges around (clockwise) bd(P). This means  $a_i = (+/-)x_j$  for some j. Consider the 1-chain  $\beta = a_1 + a_2 + ... + a_{2t} = c_1x_1 + c_2x_2 + ... + c_tx_t$  where  $c_i \in (-2, 0, 2)$ . Let the weight  $w_i$  on the edge  $x_i$  be the number of times it is traversed in the clockwise direction minus the number of times it is traversed in the counter-clockwise direction on bd(P) by the paths in U'. Consider the vectors  $w = [w_1, w_2, ..., w_t]$  and  $c = [c_1, c_2, ..., c_t]$ . The following lemma serves as a main tool in our algorithm.

LEMMA 2.2 C is null-homologous if and only if w = mc for some integer m.

PROOF. We prove that C' is null-homologous if and only if w = mc. Since C and C' are homologous the lemma follows.

Let  $\gamma$  denote the sum of the oriented triangles in T' where each triangle is oriented clockwise. If w = mc, the cycle C' represents a 1-chain that is a multiple of  $\beta$ . Since  $\beta = \delta_2 \gamma$ , we have  $C' = m\beta = m\delta_2 \gamma = \delta_2(m\gamma)$  proving C' null-homologous. To prove the other direction, assume C' null-homologous. There must exist a 2-chain d such that  $C' = \delta_2 d$ . Also, C' is carried by bd(P). Any 2-chain d with  $\delta_2 d$  carried by bd(P) must be a multiple of  $\gamma$ , see [4]. Thus  $C' = \delta_2 d = \delta_2(m\gamma) = m\delta_2 \gamma = m\beta$ . This immediately implies w = mc.

# 3 Algorithm

The polygon P is computed in O(n) time. While computing P, we maintain pointers between corresponding triangles and edges of T and T'. We traverse the cycle C on T and detect the paths

<sup>&</sup>lt;sup>2</sup>bd(\*) denotes the boundary of \*

<sup>&</sup>lt;sup>3</sup>int(\*) denotes the interior of \*

 $U = \{u_1, u_2, ..., u_r\}$  on T' in O(k) time through pointers. The paths  $U' = \{u'_1, u'_2, ..., u'_r\}$  on bd(P) can easily be detected from the end points of the paths in U. Let  $j_a$  denote the number of times an edge a on bd(P) is traversed in the clockwise direction minus the number of times it is traversed in the anti-clockwise direction. Then  $w_i = j_a + j_{a'}$  where a and a' are the partnered edges with label  $x_i$ . We can count  $j_a$ s for all edges on bd(P) by traversing the paths in U'. However, the total length of the paths in U' can be  $\Omega(kn)$  since each edge on C can contribute a path of length  $\Omega(n)$  on bd(P). To avoid this complexity increase, we count  $j_a$ s as follows.

Let  $v_1, v_2, ..., v_{2t}$  be the sequence of vertices around bd(P) in the clockwise direction. In what follows all operations on subscripts are done  $modulo\ 2t$ . Let  $a=v_iv_{i+1}$  for some  $i\in\{1,2,...,2t\}$  which is traversed in clockwise direction from  $v_i$  to  $v_{i+1}$ . To count the number of paths in U' traversing a in the clockwise direction, we have to count the number of paths starting from  $v_i$  and the number of paths that have  $v_i$  inside. Let  $\ell_{v_{i-1}}$  be the number of paths that traverse the edge  $v_{i-1}v_i$  from  $v_{i-1}$  to  $v_i$ . Then the number of paths traversing a in the clockwise direction is  $\ell_{v_i} = \ell_{v_{i-1}} + o_{v_i} - i_{v_i}$ , where  $o_{v_i}$  is the number of paths starting at  $v_i$  and  $i_{v_i}$  is the number of paths ending at  $v_i$ . To compute  $\ell_{v_i}$ , it is enough to know  $o_{v_i}$  and  $i_{v_i}$  for each vertex  $v_i$  and  $\ell_{v_1}$  for the vertex  $v_1$ . Computing  $o_{v_i}$  and  $i_{v_i}$  is straightforward. For each end vertex of the paths in U (equivalently in U') we count the number of paths starting from that vertex and the number of paths ending at that vertex. For all other vertices  $o_{v_i}$  and  $i_{v_i}$  are zero. To compute  $\ell_{v_1}$ , we number the vertices  $v_1, v_2, ..., v_{2t}$  with integer indices 1, 2, ..., 2t. Now it is simple to detect the paths in U' that include  $v_1$  in between. For this, we check in constant time if 1 is in between the integer indices of the two end vertices of a path according to the circular sequence 1, 2, ..., t, 1. The number of such paths added with  $o_{v_1}$  gives  $\ell_{v_1}$ .

Once we have computed  $\ell_{v_i}$  for each vertex  $v_i$ , it is straightforward to compute the vector w. We declare C to be null-homologous if and only if w = mc for some integer m (Lemma 2.2). Computing  $o_{v_i}, i_{v_i}$  and  $\ell_{v_1}$  while going around C (equivalently over the paths in U) takes O(k) time if C has the length k. Computing  $\ell_{v_i}$ 's while moving around bd(P) takes O(t) time. The vector c can be precomputed from P in O(t) time. Checking if w = mc takes at most O(t) time. Since t = O(n) we get an O(n + k) time complexity algorithm.

THEOREM 3.1 Let T be a triangulation of a compact 2-manifold of genus g. Given a cycle C of length k on T, there exists an O(n+k) optimal algorithm that detects if C is null-homologous.

# 4 On-line Queries

In an on-line setting, we are supposed to have a preprocessing step and then answer the null-homology of the query cycles which come on-line. For this we use the property of another invariant group of topological spaces called the fundamental group. Let P be the polygonal schema of S as constructed in Lemma 2.1 and G be the graph obtained by appropriately identifying edges of bd(P). We construct a spanning tree Y of G. Let  $b_1, b_2, ..., b_\ell$  be the edges of G not in Y and  $b'_1, b'_2, ..., b'_\ell$  be the signed edges around bd(P) corresponding to these edges. In [1], we proved that  $b_1, b_2, ..., b_\ell$  represent the generators of the fundamental group of S with respect to the relation  $b'_1, b'_2, ..., b'_{\ell} = 1$ . Also it is proved that  $\ell = O(g)$ . One dimensional homology group of a topological space can be obtained from its fundamental group by making the group operation commutative. Therefore  $b_1, b_2, ..., b_\ell$  are also generators of  $H_1(S)$  with the relation  $b'_1 + b'_2 + ... + b'_{2\ell} = c_1b_1 + c_2b_2 + ... + c_\ell b_\ell = 1$  for  $c_i \in \{-2, 0, 2\}$ .

Let  $w_i$  be the weight of the edge  $b_i$  obtained from the paths  $u'_1, u'_2, ..., u'_r$ . The given cycle C is null-homologous if and only if  $w = [w_1, w_2, ..., w_\ell] = m[c_1, c_2, ..., c_\ell]$  for some integer m. To compute w efficiently, we create another polygon Q from P in a preprocessing step as follows. The polygon Q has  $2\ell$  edges labeled  $b'_1, b'_2, ..., b'_{2\ell}$  around it. We maintain pointers from the vertices of P to the vertices of Q as follows. Let  $b'_ib'_{i+1}$  be any two consecutive edges in the sequence  $b'_1b'_2...b'_{2'}$  and  $v_1v_2...v_s$  be the vertices between  $b_i'$  and  $b_{i+1}'$  where  $v_1$  is an endpoint of  $b_i'$  and  $v_s$  is an endpoint of  $b'_{i+1}$ . All these vertices point to the same vertex v between  $b'_i$  and  $b'_{i+1}$  in Q. The polygon Q can be thought of as the polygon P with all edges in the spanning tree Y shrunk to a single vertex. The path  $u_i'$  from  $v_1$  to  $v_2$  on bd(P) is represented by the path  $u_i''$  from  $v_1'$  to  $v_2'$  on bd(Q), where  $v_1', v_2'$ are the vertices on bd(Q) corresponding to the vertices  $v_1, v_2$  respectively. The vertices  $v_1', v_2'$  can be obtained from  $v_1, v_2$  in O(1) time through pointers. Applying similar technique of section 3 on bd(Q) we can determine  $w_i$ s in O(g) time. The preprocessing step to create P and Q takes at most O(n) time altogether. Given any query cycle C on line, the end vertices of  $u_i'$  and hence  $u_i''$  can be detected in O(k) time. This is followed with an  $O(\ell) = O(g)$  step to compute  $w_i$ s and checking if  $[w_1, w_2, ..., w_r] = m[c_1, c_2, ..., c_\ell]$ . Hence on-line queries can be processed in O(q+k) time once we have an O(n) time preprocessing step.

#### 5 Conclusions

We have presented an optimal algorithm to detect null-homologous cycles on triangulated 2-manifolds. We have also given an efficient solution for the on-line version of the problem. However, this algorithm is not optimal for g > k. Finding an optimal O(k) algorithm for this problem remains open. A challenging problem is to dectect null-homologous cycles of arbitrary dimensions on arbitrary complexes. An efficient solution to this problem provides an efficient method to compute homology groups of simplicial complexes [2].

## References

- [1] T. K. Dey. A new technique to compute polygonal schema for 2-manifolds and null-homotopy detection. (1993), Manuscript.
- [2] H. Edelsbrunner and C. Delfinado. An incremental algorithm for betti numbers of simplicial complexes. Ninth Ann. Sympos. Comput. Geom. (1993).
- [3] W. S. Massey. Algebraic Topology: An Introduction. Springer-Verlag. New York (1967).
- [4] J. R. Munkres. Elements of Algebraic Topology. Addison-Wesley Publishing (1984).
- [5] H. Schipper. Determining contractibility of curves. Proc. 8th ACM Sympos. on Comput. Geom. (1992), 358-367.
- [6] J. S. Snoeyink. Topological approaches in computational geometry. *Tech. Report STAN-CS-90-1331* (1990).
- [7] G. Vegter and C. K. Yap. Computational complexity of combinatorial surfaces. Proc. 6th ACM Sympos. on Comput. Geom. (1990), 102-111.